

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/331411139>

# Evaluation and statistical optimization of a method for methylated cell-free fetal DNA extraction from maternal plasma

Article in *Journal of Assisted Reproduction and Genetics* · February 2019

DOI: 10.1007/s10815-019-01425-w

CITATIONS

0

READS

55

10 authors, including:



**Mostafa Akbariqomi**

Tehran University of Medical Sciences

16 PUBLICATIONS 21 CITATIONS

[SEE PROFILE](#)



**Reza Heidari**

Tehran University of Medical Sciences

28 PUBLICATIONS 29 CITATIONS

[SEE PROFILE](#)



**mir davood Omrani**

Shahid Beheshti University of Medical Sciences

181 PUBLICATIONS 715 CITATIONS

[SEE PROFILE](#)



**Garshasb Rigi**

Shahrekord University

68 PUBLICATIONS 80 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Project

Estimating the amount of secreted recombinant protein A using Fussy Regression insight into; determination of the optimal production conditions [View project](#)



Project

Breast cancer-VDR and lncRNA [View project](#)



# Evaluation and statistical optimization of a method for methylated cell-free fetal DNA extraction from maternal plasma

Mostafa Akbariqomi<sup>1</sup> · Reza Heidari<sup>1</sup> · Soraya Saleh Gargari<sup>2</sup> · Mir Davood Omrani<sup>3</sup> · Garshasb Rigi<sup>4,5</sup> · Nafiseh Sadat Sanikhani<sup>1</sup> · Hamid Kooshki<sup>6</sup> · Fatemeh Mahmoudian<sup>1</sup> · Mohammad Ali Mazlomi<sup>7</sup> · Gholamreza Tavoosidana<sup>1</sup>

Received: 4 October 2018 / Accepted: 18 February 2019  
© Springer Science+Business Media, LLC, part of Springer Nature 2019

## Abstract

**Purpose** Methylated cell-free fetal DNA (cffDNA) in maternal plasma can potentially be used as a biomarker for accurate noninvasive prenatal testing (NIPT) of fetal disorders. Recovery and purification of cffDNA are key steps for downstream applications. In this study, we aimed to develop and evaluate different aspects of an optimized method and compared its efficiency with common methods used for extraction of methylated cffDNA.

**Methods** Single factor experiments, Plackett-Burman (PB) design, and response surface methodology (RSM) were conducted for conventional Triton/Heat/Phenol (cTHP) method optimization. The total cell-free DNA (cfDNA) was extracted from pooled maternal plasma using the optimized method called the Triton/Heat/Phenol/Glycogen (THPG), cTHP method, a column-based kit, and a magnetic bead-based kit. In the next step, methylated cfDNA from the extracted total cfDNA was enriched using a methylated DNA immunoprecipitation (MeDIP) kit. Real-time quantitative polymerase chain reaction was performed on the RASSF1 gene and hyper region to determine the genomic equivalents per milliliter (GEq/ml) values of the methylated cfDNA and cffDNA, respectively.

**Results** The optimum values of the significant factors affecting cfDNA extraction from 200 µl of plasma were 3% SDS, 1% Triton X-100, 0.9 µg/µl glycogen, and 0.3 M sodium acetate. The GEq/ml values of methylated cffDNA extracted using the THPG method were significantly higher than for the tested extraction methods ( $p < 0.001$ ).

Mostafa Akbariqomi and Reza Heidari contributed equally to this work.

✉ Gholamreza Tavoosidana  
g-tavoosi@tums.ac.ir

Mostafa Akbariqomi  
m-akbariq@razi.tums.ac.ir

Reza Heidari  
r-heidari@razi.tums.ac.ir

Soraya Saleh Gargari  
soraya\_saleh2000@yahoo.co.uk

Mir Davood Omrani  
davood\_omrani@sbmu.ac.ir

Garshasb Rigi  
garshasbbiotech@gmail.com

Nafiseh Sadat Sanikhani  
Sanikhani@yahoo.com

Hamid Kooshki  
hmdkooshki@gmail.com

Fatemeh Mahmoudian  
Fmahmoudian@razi.tums.ac.ir

Mohammad Ali Mazlomi  
m-mazlomi@tums.ac.ir

- <sup>1</sup> Department of Molecular Medicine, School of Advanced Technologies in Medicine, Tehran University of Medical Sciences, Tehran, Iran
- <sup>2</sup> Department of Gynecology & Obstetric, Shohada Tajrish Educational Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran
- <sup>3</sup> Department of Medical Genetics, Faculty of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran
- <sup>4</sup> Department of Genetics, Faculty of Basic Science, Shahrekord University, Shahrekord, Iran
- <sup>5</sup> Research Institute of Biotechnology, Shahrekord University, Shahrekord, Iran
- <sup>6</sup> Nanobiotechnology Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran
- <sup>7</sup> Department of Medical Biotechnology, School of Advanced Technologies in Medicine, Tehran University of Medical Sciences, Tehran, Iran